

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68104 A1

- (51) International Patent Classification⁷: **A61K 31/70**, 31/505 (74) Agent: **ELBING, Karen, L.**; Clark & Elbing LLP, 176 Federal Street, Boston, MA 02110-2214 (US).
- (21) International Application Number: PCT/US01/08398 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 15 March 2001 (15.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/189,727 16 March 2000 (16.03.2000) US
60/189,811 16 March 2000 (16.03.2000) US
60/189,728 16 March 2000 (16.03.2000) US
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant: **THE MCLEAN HOSPITAL CORPORATION** [US/US]; 115 Mill Street, Belmont, MA 02178 (US). Published:
— with international search report
- (72) Inventor: **RENSHAW, Perry, F.**; 6 Newton Street, Arlington, MA 02474 (US). For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/68104 A1

(54) Title: COMPOUNDS FOR THE TREATMENT OF PSYCHIATRIC OR SUBSTANCE ABUSE DISORDERS

(57) Abstract: The invention provides methods for treating or preventing psychiatric and substance abuse disorders, involving administration of a therapeutically-effective amount of a cytosine-containing or cytidine-containing compound, creatine-containing compound, adenosine-containing, or adenosine-elevating compound to a mammal.

COMPOUNDS FOR THE TREATMENT OF PSYCHIATRIC OR
SUBSTANCE ABUSE DISORDERS

5

Statement as to Federally Sponsored Research

This invention was funded, in part, by grants DA09448 and DA11321, from the National Institute on Drug Abuse, and MH-48343 and MH-53636 from the National Institute of Mental Health. The government may have certain rights in the invention.

10

Background of the Invention

This invention relates to methods for the treatment of psychiatric or substance abuse disorders.

Psychiatric and substance abuse disorders present unique complications for patients, clinicians, and care givers. These disorders are
15 difficult to diagnose unequivocally and fear of societal condemnation, as well as lack of simple and effective therapies, often results in patients who are reluctant to disclose their symptoms to health professionals, leading to adverse societal and health consequences.

Psychiatric and substance abuse disorders include alcohol and opiate
20 abuse or dependence, depression, dysthymia, and attention-deficit hyperactivity disorder, among others, and occur in people of all ages and backgrounds.

Use of substances such as alcohol and opiates often leads to addiction and dependence on these substances, causing a variety of adverse
25 consequences, including clinical toxicity, tissue damage, physical dependence and withdrawal symptoms, and an impaired ability to maintain

social and professional relationships. The etiology of substance abuse or dependence is unknown, although factors such as the user's physical characteristics (e.g., genetic predisposition, age, weight), personality, or socioeconomic class have been postulated to be determinants.

5 Depression and dysthymia are prevalent disorders that are often chronic and associated with frequent relapses and long duration of episodes. These disorders include psychosocial and physical impairment and a high suicide rate among those affected. A lifetime prevalence of approximately 17% has been widely reported, and the likelihood of
10 recurrence is more than 50% (Angst, J. Clin. Psychiatry 60 Suppl. 6:5-9, 1999). The neurological mechanisms underlying depression and dysthymia are poorly understood, with a concomitant lack in suitable pharmacological therapies for the treatment of these disorders. Current therapies often have many adverse effects and are not suitable for
15 administration to certain cohorts. For example, depression in the elderly, particularly in those in long-term care facilities, is common and is often more refractory to treatment than depression in young or middle-aged adults; however, the elderly are particularly sensitive to the common adverse effects of many antidepressant drugs, particularly the
20 anticholinergic side effects. Similarly, therapies that are suitable for administration to adults may not be suitable for children.

 Attention-deficit hyperactivity disorder (ADHD) is a highly heritable and prevalent neuropsychiatric disorder estimated to affect 6% of the school-age children in the United States. ADHD typically occurs in
25 early childhood and persists into adulthood, but is often not diagnosed until or after adolescence. Clinical hallmarks of ADHD are inattention, hyperactivity, and impulsivity, which often respond to treatment with stimulants (e.g., methylphenidate, dextroamphetamine, or magnesium

10 Simple and effective pharmacological treatments for these disorders have proven scarce to date. It would be beneficial to provide pharmacotherapies suitable for administration to all populations, including the elderly and children, for the treatment of substance abuse and psychiatric disorders.

In general, the invention features methods of treating substance abuse or psychiatric disorders, for example, alcohol or opiate abuse or dependency, unipolar depression or dysthymia, or attention-deficit hyperactivity disorder, by administering a therapeutically-effective amount of a cytidine-containing, cytosine-containing, creatine-containing, uridine-containing, adenosine-containing, or adenosine-elevating compound to a mammal. Any of the cytidine-containing, cytosine-containing, creatine-containing, uridine-containing, adenosine-containing, or adenosine-elevating compounds of the invention may be administered separately.

3

older adult. In other preferred embodiments, the CDP-choline is administered orally and the administration is chronic.

In other preferred embodiments, a brain phospholipid (e.g., lecithin) or a brain phospholipid precursor (e.g., a fatty acid or a lipid), is also
5 administered to the mammal. In other preferred embodiments, an antidepressant is also administered to the mammal.

In other preferred embodiments, the mammal has a co-morbid neurological disease, for example, post-stroke depression.

As used herein, by "alcohol" is meant a substance containing ethyl
10 alcohol.

By "opiate" is meant any preparation or derivative of opium, which is a naturally occurring substance extracted from the seed pod of a poppy plant (e.g., *Papaver somniferum*) and which contains at least one of a number of alkaloids including morphine, noscapine, codeine, papaverine,
15 or thebaine. Heroin, an illegal, highly addictive drug is processed from morphine. For the purposes of this invention, the term opiate includes opioids.

By "opioid" is meant a synthetic narcotic that resembles an opiate in action, but is not derived from opium.

20 By "abuse" is meant excessive use of a substance, particularly one that may modify body functions, such as alcohol or opiates.

By "dependency" is meant any form of behavior that indicates an altered or reduced ability to make decisions resulting, at least in part, from the use of alcohol or opiates. Representative forms of dependency
25 behavior may take the form of antisocial, inappropriate, or illegal behavior and include those behaviors directed at the desire, planning, acquiring, and use of alcohol or opiates. This term also includes the psychic craving for alcohol or an opiate that may or may not be accompanied by a

physiological dependency, as well as a state in which there is a compulsion to take alcohol or an opiate, either continuously or periodically, in order to experience its psychic effects or to avoid the discomfort of its absence.

Forms of “dependency” include habituation, that is, an emotional or
5 psychological dependence on alcohol or an opiate to obtain relief from tension and emotional discomfort; tolerance, that is, the progressive need for increasing doses to achieve and sustain a desired effect; addiction, that is, physical or physiological dependence which is beyond voluntary control; and use of alcohol or an opiate to prevent withdrawal symptoms.
10 Dependency may be influenced by a number of factors, including physical characteristics of the user (e.g., genetic predisposition, age, gender, or weight), personality, or socioeconomic class.

By “dysthymia” or “dysthymic disorder” is meant a chronically depressed mood that occurs for most of the day, more days than not, for at
15 least two years. In children and adolescents, the mood may be irritable rather than depressed, and the required minimum duration is one year. During the two year period (one year for children or adolescents), any symptom-free intervals last no longer than 2 months. During periods of depressed mood, at least two of the following additional symptoms are
20 present: poor appetite or overeating, insomnia or hypersomnia, low energy or fatigue, low self-esteem, poor concentration or difficulty making decisions, and feelings of hopelessness. The symptoms cause clinically significant distress or impairment in social, occupational (or academic), or other important areas of functioning. The diagnosis of dysthymia is not
25 made if: the individual has ever had a manic episode, a mixed episode, a hypomanic episode; has ever met the criteria for a cyclothymic disorder; the depressive symptoms occur exclusively during the course of a chronic psychotic disorder (e.g., schizophrenia); or if the disturbance is due to the

direct physiological effects of a substance or a general medical condition. After the initial two-years of dysthymic disorder, major depressive episodes may be superimposed on the dysthymic disorder ("double depression"). (Diagnostic and Statistical Manual of Mental Disorders 5 (DSM IV), American Psychiatric Press, 4th Edition, 1994).

By "unipolar depression" or "major depressive disorder" is meant a clinical course that is characterized by one or more major depressive episodes in an individual without a history of manic, mixed, or hypomanic episodes. The diagnosis of unipolar depression is not made if: manic, 10 mixed, or hypomanic episodes develop during the course of depression; if the depression is due to the direct physiological effects of a substance; if the depression is due to the direct physiological effects of a general medical condition; if the depression is due to a bereavement or other significant loss ("reactive depression"); or if the episodes are better 15 accounted for by schizoaffective disorder and are not superimposed on schizophrenia, schizophreniform disorder, delusional disorder, or psychotic disorder. If manic, mixed, or hypomanic episodes develop, then the diagnosis is changed to a bipolar disorder. Depression may be associated with chronic general medical conditions (e.g., diabetes, myocardial 20 infarction, carcinoma, stroke). Generally, unipolar depression is more severe than dysthymia.

The essential feature of a major depressive episode is a period of at least two weeks during which there is either depressed mood or loss of interest or pleasure in nearly all activities. In children and adolescents, the 25 mood may be irritable rather than sad. The episode may be a single episode or may be recurrent. The individual also experiences at least four additional symptoms drawn from a list that includes changes in appetite or weight, sleep, and psychomotor activity; decreased energy; feelings of

worthlessness or guilt; difficulty thinking, concentrating, or making decisions; or recurrent thoughts of death or suicidal ideation, plans, or attempts. Each symptom must be newly present or must have clearly worsened compared with the person's preepisode status. The symptoms
5 must persist for most of the day, nearly every day, for at least two consecutive weeks, and the episode must be accompanied by clinically significant distress or impairment in social, occupational (or academic), or other important areas of functioning. (Diagnostic and Statistical Manual of Mental Disorders (DSM IV), American Psychiatric Press, 4th Edition,
10 1994).

By "neurological disease" is meant a disease, which involves the neuronal cells of the nervous system. Specifically included are prion diseases (e.g., Creutzfeldt-Jakob disease); pathologies of the developing brain (e.g., congenital defects in amino acid metabolism, such as
15 argininosuccinicaciduria, cystathioninuria, histidinemia, homocystinuria, hyperammonemia, phenylketonuria, tyrosinemia, and fragile X syndrome); pathologies of the mature brain (e.g., neurofibromatosis, Huntington's disease, depression, amyotrophic lateral sclerosis, multiple sclerosis); conditions that strike in adulthood (e.g. Alzheimer's disease, Creutzfeldt-
20 Jakob disease, Lewy body disease, Parkinson's disease, Pick's disease); and other pathologies of the brain (e.g., brain mishaps, brain injury, coma, infections by various agents, dietary deficiencies, stroke, multiple infarct dementia, and cardiovascular accidents). By "co-morbid" or "co-morbidity" is meant a concomitant but unrelated pathology, disease, or
25 disorder. The term co-morbid usually indicates the coexistence of two or more disease processes.

By "attention-deficit hyperactivity disorder" or "ADHD" is meant a behavioral disorder characterized by a persistent and frequent pattern of

developmentally inappropriate inattention, impulsivity, and hyperactivity. Indications of ADHD include lack of motor coordination, perceptual-motor dysfunctions, EEG abnormalities, emotional lability, opposition, anxiety, aggressiveness, low frustration tolerance, poor social skills and peer
5 relationships, sleep disturbances, dysphoria, and mood swings ("Attention Deficit Disorder," The Merck Manual of Diagnosis and Therapy (17th Ed.), eds. M.H. Beers and R. Berkow, Eds., 1999, Whitehouse Station, NJ).

By "treating" is meant the medical management of a patient with the intent that a cure, amelioration, or prevention of a disease, pathological
10 condition, or disorder will result. This term includes active treatment, that is, treatment directed specifically toward improvement of a disease, pathological condition, or disorder, and also includes causal treatment, that is, treatment directed toward removal of the cause of the disease, pathological condition, or disorder. In addition, this term includes
15 palliative treatment, that is, treatment designed for the relief of symptoms rather than the curing of the disease, pathological condition, or disorder; preventive treatment, that is, treatment directed to prevention of the disease, pathological condition, or disorder; and supportive treatment, that is, treatment employed to supplement another specific therapy directed
20 toward the improvement of the disease, pathological condition, or disorder. The term "treating" also includes symptomatic treatment, that is, treatment directed toward constitutional symptoms of the disease, pathological condition, or disorder.

By "therapeutically-effective amount" is meant an amount of a
25 cytidine-containing compound, a cytosine-containing compound, a uridine-containing compound, a creatine-containing compound, an adenosine-containing compound, or an adenosine-elevating compound sufficient to produce a healing, curative, prophylactic, stabilizing, or ameliorative effect

in the treatment of attention-deficit hyperactive disorder, unipolar depression, dysthymia,, alcohol abuse, alcohol dependency, opiate abuse, or opiate dependency.

By "cytidine-containing compound" is meant any compound that includes, as a component, cytidine, CMP, CDP, CTP, dCMP, dCDP, or dCTP. Cytidine-containing compounds can include analogs of cytidine. Preferred cytidine-containing compounds include, without limitation, CDP-choline and cytidine 5'-diphosphocholine, frequently prepared as cytidine 5'-diphosphocholine [sodium salt] and also known as citicoline.

10 By "cytosine-containing compound" is meant any compound that includes, as a component, cytosine. Cytosine-containing compounds can include analogs of cytosine.

By "adenosine-containing compound" is meant any compound that includes, as a component, adenosine. Adenosine-containing compounds 15 can include analogs of adenosine.

By "adenosine-elevating compound" is meant any compound that elevates brain adenosine levels, for example, compounds which inhibit or alter adenosine transport or metabolism (e.g., dipyridamole or S-adenosylmethionine).

20 By "uridine-containing compound" is meant any compound that includes as a component, uridine or UTP. Uridine-containing compounds can include analogs of uridine, for example, triacetyl uridine.

By "creatine-containing compound" is meant any compound that includes as a component, creatine. Creatine-containing compounds can 25 include analogs of creatine.

By "phospholipid" is meant a lipid containing phosphorus, e.g., phosphatidic acids (e.g., lecithin), phosphoglycerides, sphingomyelin, and plasmalogens. By "phospholipid precursor" is meant a substance that is

built into a phospholipid during synthesis of the phospholipid, e.g., fatty acids, glycerol, or sphingosine.

By "child or adolescent" is meant an individual who has not attained complete growth and maturity. Generally, a child or adolescent is under
5 twenty-one years of age.

By "older adult" is meant an individual who is in the later stage of life. Generally, an older adult is over sixty years of age.

The present invention provides therapeutics for the treatment of substance abuse or dependencies, unipolar depression or dysthymia, and
10 ADHD. The compounds utilized herein are relatively non-toxic, and CDP-choline, uridine, and triacetyl uridine, in particular, are pharmacokinetically understood and known to be well tolerated by mammals. The present invention, therefore, provides treatments that are likely to have few adverse effects and may be administered to children and
15 adolescents, as well as the elderly, or those whose health is compromised due to existing physical conditions.

Brief Description of the Drawings

Figure 1 is a bar graph showing the relative efficacies of CDP-choline and fluoxetine.

20 **Figure 2** is a graph showing phosphorus-31 MRS data from the human brain.

Figure 3A is a T1 weighted anatomical image of the basal ganglia and thalamus, indicating regions of interest, used to sample the T2 relaxation times, for C (caudate), P (putamen), and T (thalamus).

25 **Figure 3B** is a scatter plot of individual T2 relaxation times for the right putamen of ADHD children treated with placebo and of healthy

children. The increased T2 relaxation times seen in the ADHD sample indicate diminished regional blood volume.

Figure 4A is a graph showing the association between T2-RT in right putamen and accuracy on the performance of the computerized attention task for children with ADHD on placebo (closed circles) and normal controls (open circles). As indicated there is a significant inverse linear correlation between accuracy and T2 relaxation time (higher levels of T2-RT indicate lower perfusion).

Figure 4B is a graph showing the percent change in T2-RT in the right putamen following treatment with methylphenidate in children with ADHD. Note that the degree of response is affected by the baseline level of activity. The higher the temporal scaling the greater the activity of the subject. T2-RT change values below zero indicate enhanced regional blood volume following methylphenidate administration.

Figure 5 is a schematic illustration of the molecular structure of CDP-choline.

Detailed Description of the Invention

The invention described herein features methods for the treatment of psychiatric or substance abuse disorders such as alcohol and opiate abuse or dependence, unipolar depression, dysthymia, attention-deficit hyperactivity disorder (ADHD), and their symptoms.

To this end, the invention features the use of cytidine-containing, cytosine-containing, uridine-containing, creatine-containing, adenosine-containing, and adenosine-elevating compounds to alleviate symptoms of these disorders. A preferred cytidine-containing compound is CDP-choline (also referred to as citicoline or CDP choline [sodium salt]), a preferred

adenosine-containing compound is S-adenosylmethionine (SAdMe), and a preferred uridine-containing compound is triacetyl uridine.

The cytidine-containing, cytosine-containing, uridine-containing, creatine-containing, adenosine-containing, or adenosine-elevating
5 compounds may be co-administered with other compounds that are precursors for the synthesis of brain phospholipids, e.g., fatty acids, lipids, or lecithin.

Unipolar Depression or Dysthymia

The invention provides original data regarding the efficacy of CDP-
10 choline in human trials and demonstrates that cytidine-containing and cytosine-containing compounds can be used to treat depression. CDP-choline has been found to have two important new therapeutic properties. First, CDP-choline improves brain chemistry, e.g., increases phospholipid synthesis, in healthy adults. This effect is particularly apparent in older
15 adults. Second, CDP-choline has antidepressant effects that are similar to those of fluoxetine, a widely-used drug for the treatment of depression.

Cytidine-containing and cytosine-containing compounds are particularly efficacious in treating the elderly, and these compounds are efficacious in treating depression in patients with a co-morbid neurological
20 disease (e.g., post-stroke depression). In addition, these compounds may be administered in conjunction with, and thereby work synergistically with, phospholipids (e.g., lecithin) or compounds that are precursors for the synthesis of brain phospholipids (e.g., fatty acids or lipids).

Alcohol or Opiate Abuse or Dependence

25 Phosphorus-31 magnetic resonance spectroscopy (MRS) studies indicate that persons who are dependent upon alcohol and opiates have

decreased brain levels of phospholipids. In addition, data derived from healthy older persons, indicates that chronic administration of CDP-choline is associated with neurochemical changes consistent with phospholipid synthesis. Increasing brain levels of cytosolic adenosine also provides
5 effective therapy for alcohol or opiate abuse or dependency, because energy in the form of ATP is required to support phospholipid synthesis.

Attention Deficit Hyperactivity Disorder (ADHD)

Functional magnetic resonance imaging (fMRI) experiments in children diagnosed with ADHD indicate that symptoms of hyperactivity
10 and inattention are strongly correlated with measures of blood flow within the putamen nuclei, which are strongly dopaminergic brain regions. In addition, administration of methylphenidate, a stimulant used to treat ADHD, increases blood flow in the putamen in parallel with a decrease in motor activity. ADHD symptoms may be closely tied to functional
15 abnormalities in the putamen, which is predominantly involved in the regulation of motoric behavior. Accordingly, because cytidine-containing and cytosine-containing compounds (e.g., CDP-choline) have dopaminergic activity, these compounds may be used to treat persons diagnosed with ADHD without many of the side effects associated with
20 stimulant therapies. In particular, treatments with cytidine-containing or cytosine-containing compounds are effective in treating hyperactivity in children diagnosed with ADHD.

Cytidine-Containing and Cytosine-Containing Compounds

Useful cytidine-containing or cytosine-containing compounds may
25 include any compound comprising one of the following: cytosine, cytidine, CMP, CDP, CTP, dCMP, dCDP, and dCTP. Preferred cytidine-containing

compounds include CDP-choline and cytidine 5'-diphosphocholine [sodium salt]. This list of cytidine-containing and cytosine-containing compounds is provided to illustrate, rather than to limit the invention, and the compounds described above are commercially available, for example, 5 from Sigma Chemical Company (St. Louis, MO).

CDP-choline is a naturally occurring compound that is hydrolyzed into its components of cytidine and choline in vivo. CDP-choline is synthesized from cytidine-5'-triphosphate and phosphocholine with accompanying production of inorganic pyrophosphate in a reversible 10 reaction catalyzed by the enzyme CTP:phosphocholine cytidyltransferase (Weiss, Life Sciences 56:637-660, 1995). CDP-choline is available for oral administration in a 500 mg oblong tablet. Each tablet contains 522.5 mg CDP-choline sodium, equivalent to 500 mg of CDP-choline. Matching placebo tablets are also available. The excipients contained in both active 15 and placebo tablets are talc, magnesium stearate, colloidal silicon dioxide, hydrogenated castor oil, sodium carboxy-methylcellulose, and microcrystalline cellulose. The molecular structure of CDP-choline [sodium salt] is provided in Figure 5.

Other formulations for treatment or prevention of psychiatric and 20 substance abuse disorders may take the form of a cytosine-containing or cytidine-containing compound combined with a pharmaceutically-acceptable diluent, carrier, stabilizer, or excipient.

Adenosine-Containing and Adenosine-Elevating Compounds

Adenosine-containing or adenosine-elevating compounds provide 25 useful therapies because these compounds provide the ATP needed for phospholipid synthesis. Useful adenosine-containing or adenosine-elevating compounds include, without limitation, any compound

comprising one of the following adenosine, ATP, ADP, or AMP. One preferred adenosine-containing compound is S-adenosylmethionine (SAdMe).

In addition, compounds are known that are capable of increasing
5 adenosine levels by other mechanisms. For example, adenosine uptake can be inhibited by a number of known compounds, including propentofylline (described in U.S. Patent No. 5,919,789, hereby incorporated by reference). Another known compound that inhibits adenosine uptake is EHNA.

Other useful compounds that can be used to increase brain
10 adenosine levels are those that inhibit enzymes that break down adenosine, (e.g., adenosine deaminase and adenosine kinase). Finally, administering compounds that contain adenosine or precursors of adenosine, which are released as adenosine in vivo, can also be used.

Uridine-Containing Compounds

15 Uridine and uridine-containing compounds provide useful therapies because these compounds can be converted to CTP, a rate-limiting factor in PC biosynthesis (Wurtman et al., Biochemical Pharmacology 60:989-992, 2000). Useful uridine-containing compounds include, without limitation, any compound comprising uridine, UTP, UDP, or UMP. A preferred
20 uridine-containing compound is triacetyl uridine. Uridine and uridine-containing compounds and analogs are well tolerated in humans.

Creatine-Containing Compounds

Creatine and creatine-containing compounds provide useful therapies because these compounds, by virtue of increasing brain

phospholipid levels, can raise the levels of ATP. Creatine and creatine-containing compounds are known to be well tolerated at relatively high doses in humans.

Administration

5 Conventional pharmaceutical practice is employed to provide suitable formulations or compositions for administration to patients. Oral administration is preferred, but any other appropriate route of administration may be employed, for example, parenteral, intravenous, subcutaneous, intramuscular, intracranial, intraorbital, ophthalmic, 10 intraventricular, intracapsular, intraspinal, intracisternal, intraperitoneal, intranasal, or aerosol administration. Therapeutic formulations may be in the form of liquid solutions or suspensions (as, for example, for intravenous administration); for oral administration, formulations may be in the form of liquids, tablets, or capsules; and for intranasal formulations, 15 in the form of powders, nasal drops, or aerosols.

Methods well known in the art for making formulations are described, for example, in "Remington: The Science and Practice of Pharmacy" (19th ed.) ed. A.R. Gennaro, 1995, Mack Publishing Company, Easton, PA. Formulations for parenteral administration may, for example, 20 contain excipients, sterile water, saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes.

If desired, slow release or extended release delivery systems may be utilized. Biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be 25 used to control the release of the compounds. Other potentially useful parenteral delivery systems include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes.

Formulations for inhalation may contain excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel.

5 Preferably, the compounds of the invention, such as CDP-choline, are administered at a dosage of at least 500 mg twice daily by oral administration. Orally administered CDP-choline is bioavailable, with more than 99% of CDP-choline and/or its metabolites absorbed and less than 1% excreted in feces. CDP-choline, administered either orally or
10 intravenously, is rapidly converted into the two major circulating metabolites, choline and cytidine. Major excretion routes are lung (12.9%) and urine (2.4%); the rest of the dose (83.9%) is apparently metabolized and retained in tissues.

In general, the compounds of the invention, such as CDP-choline,
15 uridine, UTP, creatine, or SAMe, are administered at a dosage appropriate to the effect to be achieved and are typically administered in unit dosage form. The dosage preferably ranges from 50 mg per day to 2000 mg per day. The exact dosage of the compound may be dependent, for example, upon the age and weight of the recipient, the route of administration, and
20 the severity and nature of the symptoms to be treated. In general, the dosage selected should be sufficient to prevent, ameliorate, or treat alcohol or opiate abuse or dependency, or one or more symptoms thereof, without producing significant toxic or undesirable side effects. As noted above, the preferred route of administration for most indications is oral.

25 In the case of CDP-choline, there have been no reported cases of overdoses. CDP-choline toxicity is largely self-limiting, ingestion of large amounts in preclinical studies shows common cholinergic symptoms (salivation, lacrimation, urination, defecation, and vomiting).

Combination with Other Therapeutics

The cytidine-containing, cytosine-containing, uridine-containing, creatine-containing, adenosine-containing, and adenosine-elevating compounds of the invention may be administered as a monotherapy, in
5 combination with each other, or in combination with other compounds for the treatment of psychiatric and substance abuse disorders, including compounds for the treatment of alcohol or opiate abuse or dependency, or other physiological or psychological conditions associated with alcohol or opiate abuse or dependency, unipolar depression and dysthymia, and
10 ADHD.

Preferably, the compounds of the invention, may be administered in conjunction with lower doses of current treatments for these disorders, including stimulants and antidepressants. For example, the compounds of the invention may be administered with phospholipids, e.g., lecithin, or
15 with brain phospholipid precursors, e.g., fatty acids or lipids, or may be administered as an adjunct to standard therapy for the treatment of psychiatric or substance abuse disorders.

In one particular example, the compound of the invention may be administered in combination with an antidepressant, anticonvulsant,
20 antianxiety, antimanic, antipsychotic, antiobsessional, sedative-hypnotic, stimulant, or anti-hypertensive medication. Examples of these medications include, but are not limited to, the antianxiety medications, alprazolam, buspirone hydrochloride, chlordiazepoxide, chlordiazepoxide hydrochloride, clorazepate dipotassium, desipramine hydrochloride,
25 diazepam, halazepam, hydroxyzine hydrochloride, hydroxyzine pamoate, lorazepam, meprobamate, oxazepam, prazepam, prochlorperazine maleate, prochlorperazine, prochlorperazine edisylate, and trimipramine maleate; the anticonvulsants, amobarbital, amobarbital sodium, carbamazepine,

chlordiazepoxide, chlordiazepoxide hydrochloride, clorazepate
dipotassium, diazepam, divalproex sodium, ethosuximide, ethotoin,
gabapentin, lamotrigine, magnesium sulfate, mephenytoin, mephobarbital,
methsuximide, paramethadione, pentobarbital sodium, phenacemide,
5 phenobarbital, phenobarbital sodium, phensuximide, phenytoin, phenytoin
sodium, primidone, secobarbital sodium, trimethadione, valproic acid, and
clonazepam; the antidepressants, amitriptyline hydrochloride, amoxapine,
bupropion hydrochloride, clomipramine hydrochloride, desipramine
hydrochloride, doxepin hydrochloride, fluoxetine, fluvoxamine,
10 imipramine hydrochloride, imipramine pamoate, isocarboxazid,
lamotrigine, maprotiline hydrochloride, nortriptyline hydrochloride,
paroxetine hydrochloride, phenelzine sulfate, protriptyline hydrochloride,
sertraline hydrochloride, tranlycypromine sulfate, trazodone hydrochloride,
trimipramine maleate, and venlafaxine hydrochloride; the antimanic
15 medications, lithium carbonate and lithium citrate; the antiobsessional
medications, fluvoxamine, and clomipramine hydrochloride; the
antipsychotic medications, acetophenazine maleate, chlorpromazine
hydrochloride, chlorprothixene, chlorprothixene hydrochloride, clozapine,
fluphenazine decanoate, fluphenazine enanthate, fluphenazine
20 hydrochloride, haloperidol decanoate, haloperidol, haloperidol lactate,
lithium carbonate, lithium citrate, loxapine hydrochloride, loxapine
succinate, mesoridazine besylate, molindone hydrochloride, perphenazine,
pimozide, prochlorperazine maleate, prochlorperazine, prochlorperazine
edisylate, promazine hydrochloride, risperidone, thioridazine, thioridazine
25 hydrochloride, thiothixene, thiothixene hydrochloride, and trifluoperazine
hydrochloride; the sedative-hypnotic medications, amobarbital,
amobarbital sodium, aprobarbital, butabarbital, chloral hydrate,
chlordiazepoxide, chlordiazepoxide hydrochloride, clorazepate

dipotassium, diazepam, diphenhydramine, estazolam, ethchlorvynol, flurazepam hydrochloride, glutethimide, hydroxyzine hydrochloride, hydroxyzine pamoate, lorazepam, methotrimeprazine hydrochloride, midazolam hydrochloride, non prescription, oxazepam, pentobarbital sodium, phenobarbital, phenobarbital sodium, quazepam, secobarbital sodium, temazepam, triazolam, and zolpidem tartrate; the stimulants, dextroamphetamine sulfate, methamphetamine hydrochloride, methylphenidate hydrochloride, and pemoline; and the anti-hypertensive, clonidine.

10 The following examples are provided for the purpose of illustrating the invention and should not be construed as limiting.

EXAMPLES

Unipolar Depression or Dysthymia

Treatment of Human Subjects with Cytidine- or Cytosine-Containing

15 Compounds

Proton and phosphorus magnetic resonance (MR) spectroscopy studies of subjects with mood disorders have characterized two patterns of altered neurochemistry associated with depression. The first pattern indicates a change (increase or decrease) in cytosolic choline, as well as increased frontal lobe phosphomonoesters, while the second pattern points to decreased brain purines (cytosolic adenosine- and cytidine-containing compounds) and decreased nucleoside triphosphates (NTP). The former results reflect altered phospholipid metabolism, while the latter results indicate changes in cerebral energetics. Although few longitudinal studies have been conducted, these altered metabolite levels appear to be mood state, rather than trait, dependent.

To assess whether chronic CDP-choline administration leads to detectable changes in lipid metabolite resonances in phosphorus-31 MR spectra, eighteen healthy subjects (mean age: 70) were administered 500 mg of an oral formulation of CDP-choline daily for a six week period.

5 From weeks 6 to 12, half of the subjects continued to receive CDP-choline and half received placebo in a double-blind fashion. The MR data demonstrated that CDP-choline treatment was associated with a significant increase in brain phosphodiesterases ($p = 0.008$), a finding that is indicative of increased phospholipid synthesis. Neuropsychological

10 testing also revealed increases in verbal fluency ($p = 0.07$), verbal learning ($p = 0.003$), visuospatial learning ($p = 0.0001$) across all subjects at week twelve. CDP-choline administration, therefore, improves measures of verbal fluency and spatial memory in healthy adults and results in increased brain phospholipid synthesis in older adults, particularly during

15 chronic administration.

In a second study, twelve depressed subjects (mean age 40) received 500 mg of an oral formulation of CDP-choline twice daily for eight weeks. With eight weeks of treatment, mean 17-item Hamilton Depression Rating Scale (HDRS) scores decreased from 21 ± 3 to 10 ± 7

20 ($p < 0.0001$). A successful response to CDP-choline was also associated with a reduction in the proton MR spectroscopic cytosolic choline resonance in the anterior cingulate cortex. Comparable data for forty-one depressed subjects participating in imaging trials and treated with open label fluoxetine, 20 mg/day for eight weeks, demonstrated reductions in

25 HDRS scores from 21 ± 4 to 11 ± 6 ($p < 0.0001$) (Figure 1). CDP-choline and fluoxetine were associated with complete responses in 6/12 (50%) and

17/41 (41%) of the subjects, respectively (Figure 1). In depressed adults, therefore, the antidepressant effects of CDP-choline were comparable to those of fluoxetine.

These data represent the first demonstration that human brain lipid
5 metabolism can be modified using pharmacological strategies, and that, particularly in older adults, treatment is associated with improved cognitive performance. These data demonstrate that therapeutic strategies, using cytosine- and cytidine-containing compounds (e.g., CDP-choline), that are aimed at reversing biochemical alterations are beneficial
10 for the treatment of depression or dysthymia.

Use of Citicoline in a Rodent Model of Depression

The effects of citicoline were examined in the forced swim test (FST), a rodent model of depression. Because citicoline is rapidly converted to cytidine and choline, their effects were also examined in the
15 FST. Citicoline did not have antidepressant effects in rats in the FST over a range of doses (50-500 mg/kg, IP) shown to have neuroprotective effects in experimental ischemia in rodents. In fact, high doses of citicoline appeared to have small pro-depressant effects in this model. Molar equivalent amounts of cytidine (23.8-238 mg/kg, IP) had significant
20 antidepressant effects in the FST, whereas molar equivalent amounts of choline (13.7-136.6 mg/kg, IP) had significant pro-depressant effects. The optimally effective dose of cytidine (238 mg/kg, IP) did not affect locomotor activity or establish conditioned rewarding effects at therapeutic concentrations.

Alcohol or Opiate Abuse or Dependence

Measurement of Brain Phospholipids

The broad component within the phosphorus-31 MR spectrum, arising from human brain phospholipids, may be measured reliably (Figure 2). Preliminary results indicate that in persons with alcohol and/or opiate dependence, the intensity of this broad phospholipid resonance is decreased by 10-15% relative to values for comparison subjects. Accordingly, therapeutic strategies that are aimed at reversing this biochemical alteration, for example, by increasing phospholipid synthesis, are beneficial for the treatment of alcohol and/or opiate dependence.

CDP-choline Administration Leads To Increased Phospholipid Synthesis

To assess whether chronic CDP-choline administration leads to detectable changes in lipid metabolite resonances in phosphorus-31 MR spectra, eighteen healthy subjects (mean age: 70) were administered 500 mg of an oral formulation of CDP-choline daily for a six week period. From weeks 6 to 12, half of the subjects continued to receive CDP-choline and half received placebo in a double-blind fashion. The MR data demonstrated that CDP-choline treatment was associated with a significant increase in brain phosphodiesterases ($p = 0.008$), a finding that is indicative of increased phospholipid synthesis. Neuropsychological testing also revealed increases in verbal fluency ($p = 0.07$), verbal learning ($p = 0.003$), visuospatial learning ($p = 0.0001$) across all subjects at week twelve. CDP-choline administration, therefore, improves measures of verbal fluency and spatial memory in healthy adults and results in increased brain phospholipid synthesis in older adults, particularly during chronic administration.

Attention Deficit Hyperactivity Disorder (ADHD)
Functional Magnetic Resonance Imaging of Children Diagnosed with
ADHD

A new fMRI procedure (T2 relaxometry or "T2-RT") was
5 developed to indirectly assess blood volume in the striatum (caudate and
putamen) of boys 6-12 years of age under steady-state conditions. Six
healthy control boys (10.2 ± 1.5 yr) and eleven boys diagnosed with
ADHD (9.3 ± 1.6 yr) served as subjects in the study to examine fMRI
differences between unmedicated healthy controls and ADHD children on
10 either placebo or the highest dose of methylphenidate. The healthy
controls were screened using structured diagnostic interview (K-SADS-E;
Orvaschel, H. & Puig-Antich, J., The schedule for affective disorders and
schizophrenia for school-age children-epidemiologic version (Kiddie-
SADS-E), University of Pittsburgh, Pittsburgh, PA, 1987), were free of
15 any major psychiatric disorder, and had no more than 3 out of 9 possible
symptoms of inattention or hyperactivity-impulsivity by DSM-IV criteria.
Children with ADHD were included if they met criteria for ADHD on
structured diagnostic interview, and had at least 6 of 9 symptoms of
inattention or hyperactivity-impulsivity. Children with ADHD took part
20 in a triple blind (parent, child, rater), randomized, placebo-controlled
study of effects of methylphenidate (0, 0.5, 0.8, 1.5 mg/kg in divided
dose) on activity, attention, and fMRI. Children with ADHD were treated
continuously for one week with placebo or a specific dose of
methylphenidate and at the end of the week were tested for drug efficacy
25 using objective measures of attention and activity and fMRI (See
Methods) within 1-3 hours of their afternoon dose. The time between
dose and testing was held constant for each subject throughout the four
treatment conditions. Activity and attention were evaluated in

unmedicated healthy controls using the same procedure as children with ADHD, and fMRI followed within the same time frame.

T2 relaxometry, a novel fMRI procedure, was used to derive steady state blood flow measures and to test for enduring medication effects.

5 Although conventional Blood Oxygenation Level Dependent (BOLD) fMRI is a valuable technique for observing dynamic brain activity changes between baseline and active conditions, thus far it has failed to provide insight into possible resting or steady-state differences in regional perfusion between groups of subjects, or to delineate effects of chronic
10 drug treatment on basal brain function. T2 relaxometry, like BOLD, hinges on the paramagnetic properties of deoxyhemoglobin. However, the mismatch between blood flow and oxygen extraction that occurs as an acute reaction to enhanced neuronal activity in BOLD does not persist under steady state conditions. Instead, regional blood flow is regulated to
15 appropriately match perfusion with ongoing metabolic demand, and deoxyhemoglobin concentration becomes constant between regions in the steady-state. Therefore, regions with greater continuous activity are perfused at a greater rate, and these regions receive, over time, a greater volume of blood and a greater number of deoxyhemoglobin molecules per
20 volume of tissue. Thus, there is an augmentation in the paramagnetic properties of the region that is detectable as a diminished T2 relaxation time.

Conventional T2-weighted images provide only a rough estimate of T2, useful for identifying areas of pathology with markedly different T2
25 properties, such as tumors. To calculate T2-RT with sufficient accuracy to be able to reliably perceive small (ca. 2%) differences in T2 of gray matter associated with functional changes in blood volume, we used fast echoplanar imaging to establish a signal intensity decay curve based on

32 sequential measures at different echo times. For each of the 32 images, a refocused spin echo was observed.

Highly accurate laboratory-based measures of activity and attention were obtained by having the children perform a computerized vigilance test while an infrared motion analysis system captured and recorded movements (see Methods). These findings were used to ascertain associations between regional measures of T2-RT and capacity to inhibit motor activity to low levels while attending to a monotonous but demanding task.

As expected, boys with ADHD on placebo did not sit as still as healthy controls during the attention tests. They spent more time moving (temporal scaling: $F_{1,14} = 9.42$, $P = 0.008$) and had less complex movement patterns (spatial scaling: $F_{1,14} = 9.68$, $P = 0.008$). On the continuous performance task (CPT), a measure of attention, children with ADHD were less accurate (92.0% vs. 97.1%; $F_{1,14} = 2.94$, $P = 0.10$), and had a more variable response latency ($F_{1,14} = 3.11$, $P < 0.10$), though these differences did not reach statistical significance in this limited sample.

Differences in the caudate and putamen regions of children with ADHD and healthy controls, as well as the change in the T2-RT in these regions in response to methylphenidate, were also studied by imaging. The thalamus was evaluated as a contrast region in which group differences or drug effects were not expected. No significant differences emerged between ADHD children on placebo and healthy controls in bilateral T2-RT measures for the caudate nucleus ($F_{1,14} = 2.80$, $P = 0.12$). In contrast, ADHD children and controls differed markedly in bilateral putamen T2-RT measures (77.9 ± 1.1 msec vs. 76.1 ± 1.1 msec; $F_{1,14} = 9.40$, $P = 0.008$). On average, T2-RT was 3.1% higher in ADHD children

than in controls in the left putamen ($F_{1,14} = 14.5$, $P = 0.002$; Figure 3B) and 1.6% higher in the right ($F_{1,14} = 2.62$, $P = 0.13$).

For healthy controls and ADHD children on placebo, there were marked and significant correlations between motor activity and T2-RT for the putamen bilaterally, but not for caudate or thalamus (Table 1A).

Temporal scaling and average time spent immobile, two measures of activity-inactivity, correlated -0.752 ($P < 0.001$) and -0.730 ($P < 0.001$), respectively with T2-RT in putamen. The complexity of the movement pattern also correlated with T2-RT in putamen ($r_s = 0.630$, $P < 0.01$).

Similarly, in unilateral analyses, all three motor activity measures correlated with T2 measures for both right and left putamen (Table 1A).

There were also robust correlations between measures of CPT performance and T2-RT in the putamen bilaterally (Table 1B). Accuracy on the CPT correlated -0.807 ($P < 0.0001$) with T2-RT, while variability (S.D.) in response latency correlated 0.652 ($P < 0.005$). These associations were observed in both right and left putamen (Table 1B, Figure 4A). In addition, there was also a significant association between accuracy on the CPT task and T2-RT for right, but not left, thalamus. As indicated in Figure 4A, there is a significant inverse linear correlation between accuracy and T2 relaxation time (higher levels of T2-RT indicate lower perfusion).

Methylphenidate exerted robust effects on attention, enhancing performance accuracy ($F_{1,10} = 5.98$, $P < 0.05$) and reducing response variability (S.D.) from 242 to 149 msec ($F_{1,10} = 14.5$, $P < 0.005$).

Methylphenidate also exerted significant effects on activity, producing a 126% increase in time spent immobile ($F_{1,10} = 5.47$, $P < 0.05$), and increasing the complexity of the movement pattern ($F_{1,10} = 5.73$, $P < 0.05$). However, drug effects on activity were strongly dependent on the

subject's unmedicated activity level. For instance, spatial complexity increased 52.6% in the 6 subjects who were objectively hyperactive (at least 25% more active than normal controls) on placebo ($F_{1,5}=13.16$, $P < 0.02$), but was unaffected ($<8\%$ increase) in the 5 ADHD children who were not ($p > 0.6$).

T2-RT in both right and left putamen were significantly altered by ongoing treatment with methylphenidate (ANCOVA: $F_{1,9} = 12.81$, $P = 0.006$), although the response was strongly tied to the subject's unmedicated activity state (Drug \times temporal scaling covariant $F_{1,9} = 11.09$, $P = 0.008$; Figure 4B). Methylphenidate failed to exert significant effects on T2-RT in thalamus ($F_{1,9} = 0.13$, $P > 0.7$). A trend-level difference was observed in the right caudate ($F_{1,9} = 3.85$ $P = 0.08$).

Overall, as higher T2-RT corresponds to lower perfusion, the present findings of increased T2-RT in the putamen of children with ADHD, and the correlation between T2-RT and objective markers of disease severity, are consistent with some earlier studies. Furthermore, the present findings also suggest that a considerable proportion of the variance between subjects in degree of hyperactivity and inattention can be accounted for by T2-RT differences within the putamen alone.

In summary, boys with ADHD ($n = 11$) had higher T2 relaxation time (T2-RT) measures in putamen bilaterally than healthy controls ($n = 6$; $P = 0.008$). Relaxation times correlated with the child's capacity to sit still ($r_s = -0.75$, $P < 0.001$), and his accuracy in performing a computerized attention task ($r_s = -0.81$, $P < 0.001$). Blinded, placebo-controlled daily treatment with methylphenidate significantly altered T2-RT in the putamen of children with ADHD ($P = 0.006$), though the magnitude and direction of the effect was strongly dependent on the child's unmedicated activity state. A similar but non-significant trend was

observed in the right caudate. T2-RT measures in the thalamus did not differ significantly between groups, and were not affected by methylphenidate.

Methods

5 Assessment of Activity and Attention. Activity and attention data were collected as previously described (Teicher et al., J. Am. Acad. Child Adolesc. Psychiatry 35: 334-342, 1996). In brief, children sat in front of a computer and were evaluated using a simple GO/NO-GO CPT in which the subject responds to visual presentation of a target and withholds
10 response to a non-target stimuli that appear in the center of the screen at a fixed 2 second inertial interval (Greenberg et al., Psychopharmacol. Bull. 23: 279-282, 1987). The stimuli are simple geometric shapes that can be distinguished without right/left discrimination, and are designed to allow children with dyslexia to perform as well as normal controls. Three 5-
15 minute test sessions were recorded during a 30-minute test period while an infrared motion analysis system (Qualisys, Glastonbury, CT) recorded the movement of small reflective markers attached to the head, shoulder, elbow, and back of the child. The motion analysis system stored the precise vertical and horizontal position of the centroid of each marker
20 times per second to a resolution of 0.04 mm.

Results were analyzed using the concept of "micro-events." A new micro-event begins when the marker moves 1.0 millimeters or more from its most recent resting location, and is defined by its position and duration. The spatial scaling exponent is a measure of the spatial complexity of the
25 movement path, and is calculated from the logarithmic rate of information decay at progressively lower levels of resolution. The temporal scaling exponent is a scale invariant stochastic measure of percent time active.

Values range from 0 (immobility) to 1 (incessant activity), and are calculated from the slope of the log-log relationship between the duration of micro-events and their frequency (Paulus et al., Neuropsychopharmacology 7:15-31, 1992). Software for presenting stimuli, recording activity, and analyzing results was written by M. Teicher and licensed to Cygnex Inc (jtaylor@cygnex.com).

T2 Relaxometry fMRI Procedure and Relaxation Time

Computations. Children were positioned in the scanner and instructed to remain as still as possible. Images were acquired using a 1.5-T magnetic resonance scanner (Signa, General Electric Medical Systems, Milwaukee, WI) equipped with a whole-body, resonant gradient set capable of echo planar imaging (Advanced NMR Systems, Inc., Wilmington, MA), and a standard quadrature head coil for image detection. During each examination, 3 categories of images were obtained: (1) Scout images (typically T1-weighted sagittal images); (2) High resolution T1-weighted matched axial images through the ten planes for which maps of T2 were generated; and (3) 32 spin echo, echoplanar image sets, with TE incremented by 4 msec in each consecutive image set (e.g., TE (1) = 32 msec, TE (2) = 36 msec, . . . TE (32) = 160 msec) through the same ten axial planes (TR = 10 sec, Slice thickness = 7 mm with a 3 mm skip, in-plane resolution = 3.125 mm \times 3.125 mm, FOV = 200 mm). The 32 TE-stepped images were then transferred to an off-line workstation and corrected for in plane motion using a modification of the DART image registration algorithm (Maas et al., Magn. Reson. Med. 37:131-139, 1997). The value of T2-RT was then estimated on a pixel-wise basis by linear regression of the signal intensity $S(x,y,n)$ assuming an exponential decay of $S(x,y,n)$ with time constant T2-RT(x,y), such that $1/n$

$S(x,y,TE(n)) = \ln S(x,y,TE = 0) - (TE(n)/T2-RT(x,y))$, where (x,y) is the pixel position and $TE(n)$ is the spin-echo time corresponding to the n th image of the series.

Calculations of regional T2-RT were made for left and right
 5 anterior caudate, putamen, and thalamus (as a contrast region) using
 anatomic boundaries observed in T1 weighted images and conservatively
 circumscribed to avoid encroaching into ventricular space (see Figure 3A
 for regions of interest). Delineation of regions and analysis of imaging
 data was performed on coded images, and the responsible researcher was
 10 blind to the identity, diagnosis, or treatment condition of the subject. T2-
 RT was calculated from the median value of all the designated pixels, as
 the median provides a regional estimate less susceptible to contamination
 by spurious values from bordering white matter and cerebrospinal fluid
 regions than the mean.

15 The intrinsic reliability of the T2-RT measure was determined
 using a within subject procedure with head repositioning when necessary.
 There was a lag between end of the first session and start of the second
 session of ca. 5 minutes. Based on 8 within-session comparisons with
 normal adult volunteers we observed a correlation of 0.942, and an
 20 average mean value difference of -0.17% for T2-RT of the putamen.

Statistical Analyses. Differences between groups was assessed
 using ANCOVA with age as a covariate. Although the groups did not
 differ significantly in age, the behavioral and fMRI measures showed age-
 dependent changes, and ANCOVA minimized this component of the error
 25 variance. Correlations were calculated using Spearman Rank-Order test.
 Differences between behavioral and fMRI measures of ADHD subjects on
 methylphenidate vs. placebo were assessed using repeated measure

ANCOVA with placebo activity (temporal scaling) as a covariate. This was crucial in the analysis, as methylphenidate effects are strongly rate-dependent, and basal activity on placebo accounted for ca. 50% of the magnitude of the medication effect.

5 Other Embodiments

All publications and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each independent publication or patent application was specifically and individually indicated to be incorporated by reference.

10 While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure
15 that come within known or customary practice within the art to which the invention pertains and may be applied to the essential features hereinbefore set forth, and follows in the scope of the appended claims.

Other embodiments are within the appended claims.

Table 1a. Spearman Correlation between T2 relaxation time and motor behavior during CPT.

Regions	Rest-Activity Measures		Complexity Measure
	Temporal Scaling	% Time Immobile	
Bilateral			
Caudate	-0.098	-0.159	0.064
Putamen	-0.752§	-0.73§	0.63‡
Thalamus	0.152	0.194	-0.235
Unilateral			
R. caudate	-0.115	-0.115	0.196
L. caudate	-0.199	-0.270	0.054
R. putamen	-0.77§	-0.748§	0.618‡
L. putamen	-0.691‡	-0.634‡	0.534‡
R. thalamus	-0.361	0.087	0.029
L. thalamus	0.306	0.270	-0.260

‡ p = 0.055; † p < 0.05; § p < 0.005; § p < 0.001.

Table 1b. Spearman Correlation between T2 relaxation time and CPT performance.

Region	Accuracy	Response SD
<i>Bilateral</i>		
Caudate	-0.131	0.027
Putamen	-0.807§	0.652*
Thalamus	0.281	-0.135
<i>Unilateral</i>		
R. caudate	-0.020	-0.048
L. caudate	-0.357	0.087
R. putamen	-0.734§	0.629*
L. putamen	-0.708*	0.538†
R. thalamus	0.200	-0.072
L. thalamus	0.366	-0.161

† $p < 0.10$; ‡ $p < 0.05$; * $p < 0.01$; § $p < 0.001$

Claims

1. A method of treating alcohol abuse or dependency, comprising administering to a mammal a therapeutically-effective amount of a compound selected from the group consisting of a cytidine-containing
5 compound, a cytosine-containing compound, a uridine-containing compound, a creatine-containing compound, an adenosine-containing compound, and an adenosine-elevating compound.
2. A method of treating opiate abuse or dependency, comprising administering to a mammal a therapeutically-effective amount of a
10 compound selected from the group consisting of a cytidine-containing compound, a cytosine-containing compound, a uridine-containing compound, a creatine-containing compound, an adenosine-containing compound, and an adenosine-elevating compound.
3. A method of treating unipolar depression or dysthymia,
15 comprising administering to a mammal a therapeutically-effective amount of a compound selected from the group consisting of a cytidine-containing compound, a cytosine-containing compound, a uridine-containing compound, a creatine-containing compound, an adenosine-containing compound, and an adenosine-elevating compound.
- 20 4. A method of treating attention-deficit hyperactivity disorder, comprising administering to a mammal a therapeutically-effective amount of a compound selected from the group consisting of a cytidine-containing compound, a cytosine-containing compound, a uridine-containing compound, a creatine-containing compound, an adenosine-containing
25 compound, and an adenosine-elevating compound.

5. The method of claim 1, 2, 3, or 4, wherein said cytidine-containing compound is cytidine.

6. The method of claim 1, 2, 3, or 4, wherein said cytidine-containing compound further comprises choline.

5 7. The method of claim 1, 2, 3, or 4, wherein said cytidine-containing compound is CDP-choline.

8. The method of claim 1, 2, 3, or 4, wherein said cytidine-containing compound is CDP.

9. The method of claim 7, wherein said CDP-choline is
10 administered orally.

10. The method of claim 1, 2, 3, or 4, wherein said administering is chronic.

11. The method of claim 1, 2, 3, or 4, wherein said mammal is a human.

15 12. The method of claim 11, wherein said human is a child or adolescent

13. The method of claim 11, wherein said human is an older adult.

14. The method of claim 1, 2, 3, or 4, said method further comprising administering to said mammal a brain phospholipid or a brain phospholipid precursor.

15. The method of claim 14, wherein said phospholipid precursor
5 is a fatty acid.

16. The method of claim 14, wherein said phospholipid precursor is a lipid.

17. The method of claim 14, wherein said phospholipid is lecithin.

18. The method of claim 3, said method further comprising
10 administering to said mammal an antidepressant.


19. The method of claim 3, wherein said mammal has a co-morbid neurological disease.

20. The method of claim 19, wherein said co-morbid neurological disease is post-stroke depression.

1/5

FIG. 1


Open Label Response (50% reduction in HAMD-17; 8 week HAMD-17<7)
Fluoxetine, 20mg/day 18/41 = 0.44
CDP-Choline, 500mg BID 6/12=0.50



2/5

FIG. 2

Typical Decoupled P31 MRS from 5mm Axial Slices



3/5

FIG. 3B




FIG. 4A






FIG. 4B



4/5

5/5

FIG. 5



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/08398

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 31/70, 31/505

US CL : 514/49, 50, 274, 812

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/49, 50, 274, 812

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,958,896 A (RENSHAW et al) 28 September 1999, see entire pages.	2, 5-13, 19-20
X	WO 00/06174 (MASSACHUSETTS INSTITUTE OF TECHNOLOGY) 10 February 2000, see entire documents, especially page 1-4.	1-20
A,P	US 6,153,653 A (SHASHOUA) 28 November 2000, see entire documents.	19-20

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A document defining the general state of the art which is not considered to be of particular relevance	*X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*B earlier document published on or after the international filing date	*Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z document member of the same patent family
*O document referring to an oral disclosure, use, exhibition or other means	
*P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

01 MAY 2001

Date of mailing of the international search report

06 JUN 2001

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BRIAN KWON
Telephone No. (703) 308-0196

JOYCE BRIDGERS
PARALEGAL SPECIALIST
CHEMICAL MATRIX

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/08398

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN ONLINE

search terms: cytidine, cytosine, uridine, creatine, adenosin, CDP-choline, citicoline, alcohol abuse or dependency, drug abuse, opiate dependency, depression, dysthymia, attention-deficit disorder, ADHD